



**World Rowing**

**Covid-19 Pandemic:**

**Advice for training in post-peak and post-pandemic periods**

1st revision, 18 October 2020

Dr. Juergen Steinacker, Ulm University Hospital, Ulm, Germany  
On behalf of the **World Rowing Sports Medicine Commission**

Juergen Steinacker (GER), Jo Hannafin (USA), Mikio Hiura (JPN), Mike Wilkinson (CAN),  
Donia Koubaa, (TUN), Piero Poli (ITA), Petra Zupet (SLO), Tomislav Smoljanovic (CRO),  
Henning Bay Nielsen (DEN), Co-opted members: Kathryn Ackerman (USA), Fiona Wilson  
(IRL)

## Table of Contents

HEALTH SITUATION CAUSED BY THE CORONAVIRUS (Sars-CoV-2 - Covid-19).....	2
GENERAL INFORMATION ABOUT THE PANDEMIC.....	3
TRANSMISSION AND SAFETY CONSIDERATIONS .....	4
INFECTION RISK, SPORT AND RETURN TO SPORT .....	7
MITIGATION OF RISK.....	8
GENERAL INFORMATION ABOUT DEALING WITH COVID-INFECTIONS.....	11

## HEALTH SITUATION CAUSED BY THE CORONAVIRUS (Sars-CoV-2 - Covid-19)

The Coronavirus pandemic continues to evolve and is in different stages in different parts of the world. Within a country there might be areas in a peak phase with high risk of transmission close to others in which the situation may be even post-pandemic, in a so called second wave or normal. Therefore, the response to the pandemic is evolving differently across the world, in terms of governmental guidelines, the severity of the impact of COVID19 on different national populations and sports.

Acknowledging the important contribution of sport to global health and with their responsibility for rower's health, World Rowing Sports Medicine Commission, after consultation with international experts, published a guidelines document on 27 February 2020 for the spreading period and updated the guideline document on 25 March 2020 for the Peak period in which daily transmission in communities is present (Table 1)<sup>4</sup>. The "Advice for post-peak and post-pandemic periods" (this document) was published 11 June 2020. This is updated with this document to reflect the evolving global health situation.

World Rowing has published the guideline document "Covid-19 Pandemic - Return to Staging Regattas - Advice for Event Organisers and Member Federations" (2020)<sup>5</sup> and participated in the design of an "Infectious Diseases Outbreak Management Tool" with five international sports federations and WHO for sporting events with a rowing specific checklist<sup>7</sup>.

The Principles of these tools requires assessing the risk level of the sport and a particular event in both quantitative and qualitative manner<sup>7</sup> with the view to implementing mitigation measure through the following steps

- Assessing disease prevalence and the risk of transmission
- Determining the public health and sport event's mitigation preparedness
- Proposing the steps to take to further mitigate and reduce the risk.

All guidelines cannot replace the priority provisions of the state governments and the locally responsible health authorities, but serve as a rowing-specific technical contribution.

This World Rowing-guidelines document deals with the **post-peak and post-pandemic periods** by addressing a few important questions:

1. What is the transmission risk of rowing?
2. How can the transmission risk of a region, county or city be assessed for a national rowing federation, rowing club, venue?
3. How can the risk be mitigated?
4. Which measures are recommended for which disease activity?

**Typical symptoms of SARS-CoV-2**

**Fever, Dry Cough, Shortness of breath, tiredness.  
Disturbed smell and taste, sore throat, congestion, headache, inflammation of the eyes – conjunctivitis, chills, muscle and joint pain, mild diarrhoea.**

## GENERAL INFORMATION ABOUT THE PANDEMIC

According to the WHO-definition for pandemic, an outbreak has different phases. WHO defines (pandemic) severity in terms of three indicators: transmissibility of a virus, seriousness of disease and impact (Table 1)<sup>6,21</sup>.

**Table 1: WHO terminology of outbreaks**<sup>6,21</sup>

Definition	Disease activity	WHO-phase	Description
LOCAL OUTBREAK	Low	1-3	Animal to human transmission in single cases and small clusters
SPREADING PERIOD	Moderate	4	Human to human transmission in clusters and communities, disease is progressing to likely be declared pandemic
PEAK PERIOD	High	5	Significant daily community transmission with increasing new cases daily in a country
	Extra-ordinary	6	
POST PEAK PERIOD	Moderate	4	Levels of pandemic virus have dropped due to containment below peak levels with sustained decrease in community transmission
POST PANDEMIC PERIOD	low - None	1-3	Levels of virus activity have returned to low (normal) levels, but a new wave can be expected

The grading system was developed by WHO for influenza pandemic based on previous collected data<sup>2,3,19,21</sup>. Because COVID-19 is a new disease and SARS-CoV-2 a new virus, analogies to influenza, SARS-CoV-1 and other easily transmittable viral disease have to be drawn. Surveillance systems differ widely from one country to another and therefore, threshold values are difficult to compare and it is up to the state health authorities to determine and explain the level of disease activity<sup>19,21</sup>. Therefore, World Rowing strongly encourages each NF to guide their rowers on how to avoid exposure to Covid-19 by following recommendations for their regions expected to be promoted by official health authorities, such as local (government and public health) and WHO authorities.


When it can be assumed that COVID-19 disease will be prevalent for longer time, it is important to design rules and recommendations which can be adjusted also for the local and regional level. Local clusters of COVID-19-disease may arise in all phases of the pandemic<sup>2,3</sup>. *There might be areas of high prevalence (and high risk) and those of low*

*prevalence (and low risk) co-existing in regions and countries at the same time.* Those outbreaks will be assessed by health authorities, e.g. the likelihood of spreading and the possibility of containment, tracking of infected or quarantined people and the stress on the hospital system. Outbreaks in dormitories, nursing homes, in a factory may have high numbers, but may not have major effect on the other population in this region.

It is important that any rower, team support person and club member is aware of the importance of the recommended hygienic measures and social distancing to protect a safe environment for the sport. Rowing has shown with the European Rowing Championships in Duisburg (U23), Belgrade (U19) – September 2020 - and Poznan (senior) – October 2020 -, that single infected person or small groups did not spread to the other attendees with a proper hygiene and a mitigation concept.

### **Semi-confinement**

Sport teams can effectively reduce the risk of infection by semi-confinement which means voluntary restriction of social contacts also in private life and consequent hygienic measures<sup>4</sup>. Semi-confinement means that person entering the group should be illness-free for at least 10 days and be tested for SARS-CoV-2 when there is contact persons ill with COVID-19 or SARS-CoV-2 carriers.



*Figure 1. Pendulum of risk – the pandemic has various phases in different countries, regions and in and within cities. Therefore, the pandemic stresses careful risk assessment and appropriate risk mitigation. From Adami et al. 2020, with permission<sup>6</sup>.*

## **TRANSMISSION AND SAFETY CONSIDERATIONS**

### ***The basic rules to avoid being exposed to virus spread:***

- **Wash your hands frequently for a minimum of thirty seconds with soap and water or disinfected with alcoholic disinfectant when water is not available.**
- **Don't touch your mouth or nose or eyes with your hands.**
- **Practice excellent cough and respiratory hygiene.**
- **Practice physical distancing and keep distance between people.**
- **Avoid mass gatherings** where physical distancing is not possible or where local health care resources may be overwhelmed.
- **Masks covering nose and mouth** decrease the risk of transmission significantly.

### **General safety considerations for boat rowing** (Detailed phase-related considerations follow)

- As an outdoor and non-contact sport, rowing can be relatively safe due to air flow and distances possible between rowers when rowing.
- Rowing in a single is considered generally as transmission risk free.
- In crew boats, a minimum distance of 1.30m is maintained between crew members and transmission risk is minimal.
- Clubs are advised to maintain a log of crews training together so that, in case of infections, people exposed to virus can be traced.
- The virus has been shown to survive temporarily in fresh water, but transmission through fresh water is unknown<sup>30,31</sup>. World Rowing water quality guidelines apply.
- The recommendations explicitly apply to recreational and competitive rowing.
- Each rower should adopt the preventive measures, as Covid-19 follows classical dissemination via respiratory droplets and may survive on contaminated surfaces up to a few days.

### **Transmission of SARS-CoV-2 by air**

SARS-CoV-2 is one of the representatives of the species SARS-associated coronavirus, it is an enveloped RNA virus with an unsegmented genome (monopartite), i.e. only a single nucleic acid molecule (here RNA) surrounded by a capsid. It is sensitive to disinfectants and physical procedures such as heating and drying<sup>3</sup>.

The main transmission of SARS-CoV-2 is via fluid particles that are produced by coughing and sneezing and are absorbed by the opposite person via the mucous membranes of the nose, mouth and possibly the conjunctiva; contact infections are also possible. Infected fluid particles are produced by breathing, coughing, speaking and sneezing<sup>11,12</sup>. Depending on particle size, a distinction is made between droplets (larger than 5 µm) and smaller particles (infectious aerosols, smaller than 5 µm).

In principle, the likelihood of exposure to droplets and aerosols is increased within a radius of 1-2 m around an infected person. Droplets sink to the ground very quickly, aerosols can float in the air for a longer time and spread in closed rooms. The size of the particles, the air temperature, humidity and air currents are decisive factors in particle dispersion.

Virus transmission by aerosols has been documented. In indoor conditions an effective air exchange system can reduce the aerosol concentration in a room and most likely decreases infection rate. The role of aerosols in poorly ventilated rooms may be important, particularly in enclosed working areas like mines, meat industry plants, goods packing and distribution centers, but also in restaurants, clubs and bars. Sports has to consider aerosol transmission and virus spread in indoor training facilities with attention to ventilation and ability for maintaining a safe distance.

**Table 2: Summary of pandemic phases of Sars-CoV-2 – Covid-19, safety levels, distancing and precautions (for details see below)**

Phase	Prevalence	Safety level	WHO-recommended Distance (m)	Rowing	Boat house	Weights room	Ergo exercise
LOCAL OUTBREAKS	Single cases and small clusters	Level 1 Normal sports <b>Hygiene rules always apply</b>	No regulation	Crew	open	Open Disinfection Hygiene	Open
SPREADING PERIOD	In clusters and first communities	Level 2 Sport with safety precautions	1.0	Crew <i>with safety</i>	Limited access	Hygiene	Hygiene
Pandemic phase / ACTIVE OUTBREAK	High community prevalence	Level 3 Sport restricted	1.5	Single	closed	Closed	Private
Post Peak phase / OUTBREAK UNDER CONTROL	Low community prevalence – widely contained	Level 2 Sport with safety precautions	1.0	Crew <i>with safety</i> <sup>1</sup>	Limited access	5 rowers + 1 coach <sup>a</sup> Disinfection <sup>b</sup> and Enhanced cleaning	1.5 m distance between ergometers Good ventilation
CONTAINED OUTBREAK - Post Pandemic phase	Controlled / contained	Level 1 Normal sports <b>Hygiene rules always apply</b>	No regulation	Crew	open	Open Disinfection Hygiene	Open

- a. Number of crew members or group size depends on local prevalence and conditions and on assessment by public health authorities
- b. Disinfection of each equipment contact point before and after use

World Rowing's focus is to provide advice on measures which effectively prevent exposure as well as measures that limit the environmental risk and the risk in the social environment. Furthermore, risk is greatly reduced by the fact that the rowing groups are relatively small and training groups and team do not change much within time. Rowers also come from socially very reliable groups with a high level of health awareness and very good hygiene behavior. In addition, chains of infection can be traced very well and SARS-CoV-2-transmission in rowing can be effectively controlled.

**The risk of transmission can be mitigated significantly by masks** covering nose and mouth<sup>10,11,13</sup>. Masks can decrease the number of aerosols and particles inhaled significantly and therefore decrease the likelihood of infection or the severity of disease, because viral load is decreased. Masks should be used consequently outside the rowing boat, particularly, in boat house facilities, during meetings and travel<sup>13,27</sup>. Face shields are not effective, they only protect against direct exposure by droplets during coughing and speaking, but not protect against aerosols and should not be used<sup>10,11,13</sup>.

Exercise capacity is decreased by wearing a mask<sup>10,14</sup>. During ergometer training, face masks may interfere with training but rowers training on an erg exhale aerosol. There is no data on aerosols during ergometer rowing, but it is recommended to increase distance between ergometers and to ensure good ventilation.

## INFECTION RISK, SPORT AND RETURN TO SPORT

### **Return to Training Considerations in post-peak and post-pandemic periods:**

- What is the rate of community COVID-19 transmission, and therefore the risk of infection, low enough to allow group training or sharing of equipment?
- Are rowers and staff trained sufficiently in hygienic measures and responsible practices?
- Are there appropriate resources (hand washing facilities, soap, disinfectant, face masks) and are hygienic measures secured (cleaning of facilities, surfaces, toilets, ...)

WHO recommends to evaluate the (pandemic) severity in terms of three indicators: transmissibility of a virus, seriousness of disease and impact in the so-called PISA-assessment (Pandemic Influenza Severity Assessment) which is used in analogy also for COVID-19<sup>21</sup>. The Cut-off Points by method for threshold settings for transmissibility and impact are defined by epidemiological methods. The seriousness of disease will be assessed by official health authorities and the impact is related to the use of emergency capacities and hospital beds of the country<sup>21</sup>.

The rate of acute respiratory illnesses (ARI) surveillance is established for analysis of transmissibility of a virus according to the PISA-method<sup>21</sup>. For the comparability of transmissibility among countries and seasons the moving epidemic method (new cases in the last seven days per 100.000 inhabitants) is widely used in Europe<sup>15,16,17</sup> and also a recommended method by WHO<sup>21</sup>.

Cut-off points for threshold settings for transmissibility are used as recommended by the German Robert-Koch-Institute<sup>17</sup> with thresholds of lower than 50/100.000 per inhabitants for moderate risk and 20/100.000 per inhabitants for low risk. These cut-points allow to assess and compare the risk of transmission for COVID-19 between cities, regions and countries and allow organizers to evaluate the risk and tailor their mitigation procedures to the local or regional context (Table 3)<sup>2,3,21</sup>.

Health authorities should adjust and tailor their approach to the local context<sup>2,3,21</sup>. Health authorities may also use more variables than transmissibility to assess the risk like regional intensive care use or local clustering.

When new clusters of Sars-CoV-2 - Covid-19 appear in a region, the relevance for this population and the risk of transmission has to be assessed by local health authorities<sup>21</sup>. When an outbreak is contained (e.g., in a work place, nursing home ...) the risk for the other population may be much lower.

**Table 3: Risk attribution and mitigation procedures according to different regional scenarios of Sars-CoV-2 - Covid-19 according to the WHO-guidelines<sup>2,3, 9, 21</sup> and using the RKI-threshold cut-off for transmissibility<sup>17</sup>**

Risk	Scenario	Region / County: New Cases per 100.000 inhabitants in 7 days <sup>17</sup>	WHO- recommended Distance (m)	Rowing	Boat house	Weights room	Ergo exercise
High & extraordinary	Countries / Regions experiencing larger outbreaks of local transmission (Community transmission).	More than 50 cases per 100,000 in 7 days	1.5	Single	Closed	Closed	Private
moderate	Countries / Regions experiencing cases clusters in time, geographic location, or common exposure (Contained clusters of cases);	Between 20 and 50 cases per 100,000 in 7 days, or, higher prevalence but contained clusters	1.0	Crew <i>with safety</i>	Limited access	5 rowers + 1 coach <sup>a</sup> Disinfection <sup>b</sup> Enhanced cleaning	1.5 m distance between ergometers Good ventilation
Low	Countries / Regions with one or more cases, imported or locally detected (Sporadic cases);	Less than 20 cases per 100,000 in 7 days	1.0	Crew <sup>c</sup> <i>with safety</i>	Limited access	10 rowers + 1 coach <sup>1</sup> Disinfection <sup>b</sup> Hygiene	Good ventilation
No	Countries / Regions with no cases (No cases);	Not applicable	No regulation	Crew	Open	Open Disinfection Hygiene	Open

- a. Number of crew members or group size depends on local prevalence and conditions and on assessment by public health authorities
- b. Disinfection of each equipment contact point before and after use
- c. Crew size can be increased to big boats due to local prevalence and conditions

## MITIGATION OF RISK

### 1. Minimize risks in all areas (at all times)

- a. **Rowers and support staff must not participate in training if they show any symptoms of illness or have been exposed to a COVID-positive person.** They must stay at home or in isolation and call their family doctor and follow their instructions; this also applies to accompanying persons. The training group or other contacts must be informed immediately. The use of common facilities and training rooms for ill rowers is not permitted.
- b. **Training is not allowed if there are symptoms of illness.**
- c. Typical symptoms of Sars-CoV-2 - Covid-19 illness are fever, dry cough, disturbed smell and taste, conjunctivitis, mild diarrhea, tiredness or shortness of breath.

### 2. Observe distance rules

Table 3 and local assessment by public health authorities

- a. When training outdoors alone, a person has a low risk of infection. The risk can be increased secondarily by the proximity to training partners and the length of time in that proximity. It is therefore important to keep a distance according to the risk phase (Table 1) when interacting on the boathouse premises, especially when maintaining equipment and launching the boats.
- b. Wearing of a mouth and nose mask decreases risk of transmitting the virus to others, particularly when distancing is not possible (work place or at the dock). Community masks may decrease droplets depending on material between 10 and 95% (dense cotton seems to be best), surgical masks by 85-97%. Eye protection inhibits further infection. Ensure proper training on how to handle a mask.
- c. With increased risk, rowers should wear a mask in the boat house and on the dock at all times.

### **3. Reducing the size of training groups in the post-peak period (risk medium)**

Table 3 and local assessment by public health authorities

- a. Water training is always permitted in singles.
- b. Training in a crew is possible if all members are from a household or quarantined together.
- c. Reduce the size of a training group to five persons (high risk) or 10 (medium risk) plus the coach (Table 3).
- d. Training in a crew boat may also be possible when permitted by local regulations and according to risk (Table 3).
- e. Training group should not be changed in terms of personnel. Training groups can be seen as semi-confined and of lower risk when members keep social distancing outside their training.
- f. If the respective legal regulations concerning the distance or composition of groups require it, an intermediate seat can be left free in boats. Official regulations regarding the extent of contacts are decisive for the composition of crews / training groups.
- g. If a coxswain is participating, face mask could be worn.
- d. Oar handles should be cleaned and disinfected after training. Disinfectant cleaning agents approved by health authorities are recommended.

### **4. Boathouse use**

- a. Reopening of rowing clubs and national training centers is upon discretion of the health authorities.
- b. World Rowing SMC recommends strictly limiting the use of boathouse facilities, particularly dressing rooms, showers and training facilities, depending on risk assessment. (Table 2). If possible, rowers should change and shower at home rather than at boat house facilities. When the risk is rated low, reopening should be considered (Table 3).
- c. Rooms should be ventilated regularly.
- d. Daily cleaning of all facilities and maintaining hygiene is mandatory. Washrooms, toilets (surfaces and contact points) and door handles should be disinfected frequently. Do not use fan hand dryers; use single user paper towels. Close the hood in toilets before flushing.

### **5. Exercise and weight rooms**

- a. Access to exercise and gym rooms must be strictly controlled to ensure cleanliness and hygiene. For room occupation where indoor training is permitted, a room size of 5 sqm per person, with controlled flow patterns to ensure distancing, is recommended. This should be eased when disease activity and risk is low (Table 3).
- b. Ergometer exercise should be done with open windows and a distance > 1.5 m between ergometers when disease activity and risk is moderate (Table 3).
- c. Prior to first-time use, all surfaces and the floor must be treated with an approved disinfectant cleaning agent. This must be repeated at least weekly; daily cleaning is recommended.
- d. All equipment, ergometers, dumbbells, etc. must be treated with an approved disinfectant before and after use at all contact points and exposed surfaces.
- e. Except during ergometer training, a mask should be worn. Hands must be washed before and after training.

## **6. Competition and regattas:**

- a. In the current situation, World Rowing aims to gain experience through regional and continental events and encourages all NF's to organize safe competitions to enable rowers to compete.
- b. The risk in rowing events is related to the number of persons gathered and the possible transmission by inadvertent carriers.
- c. Hygiene rules, social distancing have to be observed carefully. Face mask should be used on the venue in pandemic
- d. World Rowing has published a guidelines document for Event Organisers and Member Federations<sup>5</sup>.
- e. Pandemic-specific regulations of local health authorities have to be followed,
- f. The WHO Risk assessment tool helps to assess risks, mitigate spread, and enhance communication with local health authorities<sup>9</sup>.

## **7. Prevention:**

- a. Where air travel is necessary, face masks and eye protection should be used, but may not fully protect against infection. In the airplane, disinfection of the seat tray is recommended. Personal containers of hand sanitizer or disinfectant should be carried and used regularly.
- b. A varied, vitamin-rich diet with sufficient vegetables and fruit helps prevent infections or reduce symptoms by improving immunity. Rowers should also drink enough to keep mucous membranes moist. In cold weather, the immune function of the mucous membranes is particularly disturbed, so keep the respiratory tract as warm as possible.
- c. Food should not be served in buffet style. Wash hands with soap before and after meals.

## **8. Drawing up a hygiene plan**

- a. Clubs and event organizers are requested to draw up a hygiene plan and document the measures taken. This should include the maximum occupancy according to the risk evaluation.
- b. For events, a hygiene plan is necessary. WHO guidelines on mass gatherings and World Rowing guidelines should be considered. Local health authority regulations should also be followed.

## GENERAL INFORMATION ABOUT DEALING WITH COVID-INFECTIONS

***The basic rules for those who are infected or do feel not well:***

- **Typical symptoms of COVID-19 are fever, dry cough, disturbed smell and taste, inflammation of the eyes - conjunctivitis, mild diarrhea, but also tiredness or shortness of breath can be felt.**
- **You should stay in self-quarantine at home and communicate with your doctor.**
- **Avoid sports practice in groups.**
- **Go into self-quarantine. Do not get close to your partner, family members or other people, keeping a minimum distance of 2 meters.**
- **Eat separately and use a separate toilet.**
- **Wash your hands regularly with soap.**
- **Contact your medical doctor and others by phone.**
- **Only when cleared by your doctor can you return to your normal routine.**

### **1. If rowers feel sick**

- a. With symptoms, rowers and entourage should take a break from training and call their team or family doctor and follow their instructions. They should inform team members and other recent contacts.
- b. Such rowers and entourage should be separated from the rest of the team, self-isolate and carefully disinfect their hands. They should wear a face mask during acute symptoms and ensure that the mask is used correctly.
- c. The separation should be at least 2-3 days more than the actual signs of infection and should be decided by the attending doctor. It is wise to have separated toilets for such rowers to break the transmission of virus on hard surfaces.

### **2. How can I exclude Coronavirus (Sars-CoV-2 / COVID-19) infection if I have symptoms?**

- a. Fever is uncommon with most colds and upper respiratory infections but is more common with COVID-19 and the flu.
- b. The probability of infection with the coronavirus (Sars-CoV-2 / COVID - 19) is very high in regions where the infection is widespread (community infection). Caution is generally required for any contact outside the family.
- c. Most infections with Coronavirus (Sars-CoV-2 / COVID - 19) are mild (in many there may be very few or no symptoms), but more severe complications may arise, particularly in older and immunocompromised people.
- d. Coronavirus (Sars-CoV-2 / COVID - 19) infection should be considered a possible cause of infection in acute respiratory illness and should be assessed by a medical doctor.
- e. Transmission time appears to be 14 days. During that time an infected person may transmit the virus to others, even if there are no clinical signs of infection. During this asymptomatic period, the virus may also not be detectable by the PCR (Nasal swab) or serological (Blood) test.

### 3. Return to Sport after Coronavirus infection (SARS-CoV-2 / COVID-19)

*(NOTE: All the recommendations for return to sport after COVID-19 are expert consensus; no controlled studies available).*

- a. Rowers and entourage **who are ill and had contact** with an infected individual or someone who traveled to affected regions should seek medical advice in a public health office where Coronavirus (Sars-CoV-2 / COVID - 19) testing can be performed (following and according to local health instructions).
- b. **Work-up-plans for rowers** have been published<sup>6, 18,22,23,24</sup>
- c. All rowers with symptoms should be seen by their sports medicine doctor before resuming training.
- d. In rowers with a **positive SARS-CoV-2 test WITHOUT** signs of infection or symptoms, a 14-day-quarantine will likely be required, but it is important to follow local public health or WHO recommendations regarding further testing and isolation. Only light activity for 2 weeks and then return to training should be considered<sup>22,23,24</sup>.
- e. Rowers with a **positive SARS-CoV-2 test WITH symptoms** but WITHOUT confirmed pneumonia, should be examined by their doctor. No heavy training for 2 to 4 weeks, then gradual return to training should be considered<sup>6, 18,22,23,24</sup>.
- f. In rowers with a **positive SARS-CoV-2 test WITH pneumonia**, exercise testing with measurement of oxygen saturation should be performed. Further examinations should take place as recommended. No sports for at least 4 weeks, then gradual return to training under doctor's supervision should be considered<sup>6, 18,22,23,24</sup>.
- g. In rowers with **positive SARS-CoV-2 test WITH suspected or confirmed myocarditis** with/without pulmonary involvement, cardiac testing should be performed. If there is no evidence of myocarditis, return to sports may be permitted. If myocarditis is diagnosed, no sports for at least 3 months with gradual return to training after reevaluation by and under the supervision of a cardiologist<sup>24,25</sup>.

### 4. Is there a role for vaccination?

- a. There is no vaccination presently available for Coronavirus (Sars-CoV-2 / COVID-19). Many pharmaceutical companies continue to work on vaccines but availability and efficacy is still unknown.
- b. World Rowing SMC continues to recommend annual vaccination with quadrivalent flu vaccines against viral influenza for rowers and their entourage. However, COVID-19 is not an influenza strain and is not prevented by this vaccine. The recommendation is to prevent influenza. Flu vaccination is most effective in the northern hemisphere in November and in the southern hemisphere in June and is updated annually by the WHO.

### 5. The role of prospective SARS-CoV-2 screening?

- a. Testing before or after an event is advised in the case of positive cases or ill cases and contacts in a NF. National authorities may demand testing when crossing borders.
- b. World Rowing presently does not advise general asymptomatic testing when a very low rate of positive cases is expected. For example, when specificity of the PCR test for detection of virus antigen is assumed very high at 99.9% and prevalence of COVID is assumed to be very low at 0.1%, this would generate 50 % false positive cases, when specificity is already 99 %, it would generate 90% false positives (prevalence 0.1%).
- c. A second test significantly increases the diagnostic certainty. For groups of people who have spent a long time together (e.g. teams in training camps or families), the probability of the correctness of positive test results increases with the number of positive test results within the community (because infection within the community is likely).

- d. Conversely, the opposite is true: if there are only a few positive test results in communities, this reduces the probability that these results are correct. Individual positive tests in communities (without documented contact to risk areas or symptoms etc.) must therefore be treated with great caution. Such test results would in fact be (arithmetically) more false than correct.
- e. So if a team is tested positive in the majority of the training camp, then the results are almost certainly correct. However, if only one or two of the 30 people in the team test positive, a false positive result is more likely.
- f. The PCR-test is very specific and has a high sensitivity. There are also various rapid tests available. These have lower specificity and sensitivity which means a higher rate of false positive and negative results and therefore used and interpreted with caution. But when technology is improved, rapid test may a useful tool.

#### **Where is information available?**

**The following webpages allow the most recent international information:**

- **WHO-Websites:**
  - <https://www.olympic.org/athlete365/covid-19-resources/>
  - <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>
- **WORLD ROWING (will be updated regularly)**
  - [http://www.worldrowing.com/mm/Document/General/General/14/28/56/WORLD\\_ROWINGReturntoRegattasGuidance-Version2180920\\_Neutral.pdf](http://www.worldrowing.com/mm/Document/General/General/14/28/56/WORLD_ROWINGReturntoRegattasGuidance-Version2180920_Neutral.pdf)
- **WHO Risk assessment tool** helps to assess risks, mitigate spread, and enhance communication with local health authorities:
  - <https://www.who.int/publications-detail/key-planning-recommendations-for-mass-gatherings-in-the-context-of-the-current-covid-19-outbreak>
  - Event risk and mitigation-tool with rowing-specific mitigation checklist:  
<https://idom.worldathletics.org/>
- **Return to Sport after Covid:**
  - <https://www.germanjournalsportsmedicine.com/archiv/archive-2020/issue-5/position-stand-return-to-sport-in-the-current-coronavirus-pandemic-sars-cov-2-covid-19/>

#### **References:**

1. WHO. WHO global epidemiological surveillance standards for influenza. Geneva: World Health Organization (WHO); 2014  
[http://www.who.int/influenza/resources/documents/influenza\\_surveillance\\_manual/en/](http://www.who.int/influenza/resources/documents/influenza_surveillance_manual/en/)
2. WHO Pandemic phase descriptions and actions, WHO,2020
3. WHO Critical preparedness readiness and response actions COVID-10 2020-03-22\_FINAL-eng
4. Steinacker JM, and Sports Medicine Commission of World Rowing. Fact Sheet: Health Situation for Rowers in the Current Coronavirus Pandemic (SARSCoV-2 / COVID-19). Dtsch Z Sportmed. 2020; 71: 85-86. doi:10.5960/dzsm.2020.431
5. WORLD ROWING Sports Med Commission (2020) Covid-19 Pandemic - Return to Staging Regattas. Advice for Event Organisers and Member Federations
6. Löllgen H, Bachl N, Papadopoulou T, et al. (2020) Recommendations for return to sport during the SARS-CoV-2 pandemic. BMJ Open Sport & Exercise Medicine. 2020;6:e000858. doi:10.1136/bmjsem-2020-000858
7. Adami PE, Cianca J, McCloskey B, et al. (2020) Infectious Diseases Outbreak Management Tool for Endurance Mass Participation Sporting Events. An international Effort to Counteract the

- COVID-19 Spread in the Endurance Sport Setting. *Br J Sports Med* 2020; Epub ahead of print 0:1–2. doi:10.1136/bjsports-2020-103091
8. Muniz-Pardos, B., Shurlock, J., Debruyne, A. et al. (2020) Collateral Health Issues Derived from the Covid-19 Pandemic. *Sports Med - Open* 6, 35 (2020). <https://doi.org/10.1186/s40798-020-00267-6>
  9. WHO 2020, How to use WHO risk assessment and mitigation checklist for Mass Gatherings in the context of COVID-19. WHO-2019-nCoV-Mass\_Gatherings\_Sports-2020.1-eng.
  10. Seto WH, Tsang D, Yung RW, Ching TY, Ng TK, Ho M, Ho LM, Peiris JS; Advisors of Expert SARS group of Hospital Authority. Effectiveness of precautions against droplets and contact in prevention of nosocomial transmission of severe acute respiratory syndrome (SARS). *Lancet*. 2003 May 3;361(9368):1519-20. doi:10.1016/s0140-6736(03)13168-6
  11. Morawska L, Cao J. *Environ Int*. 2020; 139: 105730. doi:10.1016/j.envint.2020.105730
  12. Van Doremalen N, et al.. *N Engl J Med*. 2020; 382: 1564-1567. doi:10.1056/NEJMc2004973
  13. Prather KA, et al. (2020) Reducing transmission of SARS-CoV-2. *Science* 368, 1422-1424 DOI:10.1126/science.abc6197
  14. Fikenzler S et al. (2020) Effects of surgical and FFP2/N95 face masks on cardiopulmonary exercise capacity *Clin Res Cardiol* <https://doi.org/10.1007/s00392-020-01704-y>
  15. Vega T et al. Influenza surveillance in Europe: comparing intensity levels calculated using the moving epidemic method. *Influenza Other Respir Viruses*. 2015;9(5):234–246 <https://www.ncbi.nlm.nih.gov/pubmed/26031655>
  16. Vega T et al. Influenza surveillance in Europe: establishing epidemic thresholds by the moving epidemic method. *Influenza Other Respir Viruses*. 2013;7(4):546–558 <https://www.ncbi.nlm.nih.gov/pubmed/22897919>
  17. Pandemieplan – COVID-19 – neuartige Coronaviruserkrankung. Robert-Koch-Institut, Berlin, 2020
  18. Nieß et al. Position Stand: Return to Sport in the Current Coronavirus Pandemic (SARS-CoV-2 / COVID-19) *Dtsch Z Sportmed*. 2020; 71: doi:10.5960/dzsm.2020.437
  19. Overview of Public Health and Social Measures in the context of COVID-19 - WHO/2019-nCoV/PHSM\_Overview/2020.1
  20. WHO-WHE-IHM-GIP-2017.2-eng.pdf
  21. Pandemic Influenza Severity Assessment (PISA), WHO, Geneva, 2017 [https://www.who.int/influenza/surveillance\\_monitoring/pisa/guidance/en/](https://www.who.int/influenza/surveillance_monitoring/pisa/guidance/en/)
  22. The Australian Institute of Sport (AIS) Framework for Rebooting Sport in a COVID-19 Environment — Executive Summary
  23. Return to Training Considerations Post-COVID-19 United States Olympic & Paralympic Committee. April 28, 2020 – v0.11
  24. Nieß AM et al. Recommendations for exercise testing in sports medicine during the current pandemic situation (SARS-CoV-2 / COVID-19). 2020; 71: E1-E2. DOI: 10.5960/dzsm.2020.438
  25. Baggish AL et al.: the resurgence of sport in the wake of COVID-19: cardiac considerations in competitive rowers. <https://blogs.bmj.com/bjism/2020/04/24/the-resurgence-of-sport-in-the...>
  26. McCloskey B et al. Mass gathering events and reducing further global spread of COVID-19: a political and public health dilemma. *Lancet* 2020;395(10230):1096-99. doi: 10.1016/s0140-6736(20)30681-4
  27. Carmody S et al. When can professional sport recommence safely during the COVID-19 pandemic? Risk assessment and factors to consider. *Br J Sports Med* 2020 doi: 10.1136/bjsports-2020-102539
  28. Barker-Davies RM et al. The Stanford Hall consensus statement for post-COVID-19 rehabilitation. *Br J Sports Med* 2020;0:1–11. doi:10.1136/bjsports-2020-102596
  29. Wackerhage H et al.. Sport, exercise and COVID-19, the disease caused by the SARS-CoV-2 coronavirus. *Dtsch Z Sportmed*. 2020; 71: E1-E12. doi:10.5960/dzsm.2020.441
  30. CDC recommendation for aquatic venues, Bethesda, 2020. <https://www.cdc.gov/coronavirus/2019-ncov/community/parks-rec/aquatic-venues.html>
  31. WHO Water sanitation hygiene and waste management for covid-19, WHO, 2020 <https://www.who.int/publications-detail/water-sanitation-hygiene-and-waste-management-for-covid-19-2>